

Impossibility of Spin Polarized States for Neutron Star / Proto-neutron Star Matter in β -Equilibrium Condition

Sutapa Ghosh^{a)*}, Soma Mandal^{a) †} and Somenath Chakrabarty^{a),b)‡}

^{a)}*Department of Physics, University of Kalyani, Kalyani 741 235, India and* ^{b)}*Inter-University Centre for Astronomy and Astrophysics, Post Bag 4, Ganeshkhind, Pune 411 007, India*

(February 1, 2008)

PACS:97.60.Jd, 97.60.-s, 75.25.+z

Key Words: Ferro-magnetic transition, Neutron stars, Proto-neutron stars, Millisecond pulsars.

It is shown explicitly that a ferromagnetic transition of neutron star (NS) / proto-neutron star (PNS) matter in the β -equilibrium condition with $\sigma-\omega-\rho$ exchange type of mean field approximation can actually occur if and only if the neutrinos remain trapped within the system, and perhaps it is also necessary for the neutrinos / anti-neutrinos to carry some finite non-zero mass. It is further shown that the electrons also play a significant role in this transition process. It is therefore, very much unlikely that such a transition of spin polarization can really take place in old neutron stars of very low temperature, whereas, the possibility of spontaneous ferromagnetic transition cannot be ruled out in a newly born PNS.

The origin of strong magnetic field of pulsars, which are also believed to be the rotating neutron stars, is not exactly known till today [1]. There are so many models of which the idea of flux conservation during collapse gives rise to a field strength of $\approx 10^{12}$ G for some class of neutron stars. This field is known as fossil remnant from the progenitor star, with a typical magnetic field strength of $\sim 10^4$ G. The other alternative model for the generation of magnetic field is by some kind of dynamo process or electric current flowing in highly conducting material of the neutron stars after their formation [1–5]. Now the studies on magnetic field evolution of neutron stars by population synthesis codes show that the magnetic field decay time scale for typical neutron stars (not the magnetar like exotic objects) is $\sim 10^7 - 10^9$ yrs and the residual magnetic field $\sim 10^8$ G [6] (see also [7–9]). This is also the order of magnitude of magnetic field strength for the millisecond pulsars, which are believed to be re-generated and started their rotations with very high angular velocity after crossing the death line from pulsar graveyard by the accretion of matter [1], i.e., such objects are very old (and also cold) neutron stars. If this residual magnetic field is assumed to be permanent and not generated by some dynamo process (the assumption is possibly true in the case of very old neutron stars) within the star, one of the possibility of such permanent magnetic field is a spontaneous ferromagnetic transition by the spin alignment of permanent elementary magnetic dipoles of neutrons, protons etc. Since it is generally believed that such a transition may be the origin of magnetic field, in particular the remnant magnetic field of millisecond pulsars which are assumed to be very old neutron star, we have considered it to be an important issue. Of course, if a spontaneous ferro-magnetic transition is not possible in neutron star matter in β -equilibrium in presence of non-degenerate neutrinos, in spite of that we realized that this very old problem of spontaneous ferro-magnetic transition has become extremely interesting by itself (also has some academic importance). To the best of our knowledge such studies have not been done before and this is the first attempt to investigate this anomaly.

In this article we shall show that a spontaneous ferromagnetic transition is completely forbidden in a NS if the matter is in β -equilibrium and neutrinos leave the system immediately after their formation (i.e., non-degenerate). However, such a transition is possible if the produced neutrinos remain trapped within the neutron matter system (i.e. in PNS) for quite a long time at the initial stage of evolution. Now the ferromagnetic materials are paramagnetic but, because of interactions between atoms, show drastically different behavior. Below the Curie temperature, ferromagnetic substance show spontaneous magnetization, that is all the magnetic moments in a microscopically large region called a domain are aligned. The application of an external field tend to cause the domains to change and the moments in different domains to line up together leading to the saturation of the bulk magnetization. Removal of the field leaves a considerable fraction of the moments still aligned, giving a permanent magnetization. We further assume a $\sigma-\omega-\rho$ meson type of mean field interaction picture in the hadronic sector, whereas, electrons and neutrons are assumed to be non-interacting.

In a NS / PNS the dynamic nature of chemical equilibrium among the constituents is mainly dominated by the following weak interaction processes (URCA processes)

*E-Mail: sutapa@klyuniv.ernet.in

†E-Mail: soma@klyuniv.ernet.in

‡E-Mail: somenath@klyuniv.ernet.in

$$\begin{aligned}
n &\rightarrow p + e^- + \bar{\nu}_e \\
p + e^- &\rightarrow n + \nu_e
\end{aligned} \tag{1}$$

Now in a typical neutron star, considering the spontaneously spin polarization of hadronic matter, in which neutrinos leave the system as soon as they are produced, the conditions for dynamical β -equilibrium is given by

$$\begin{aligned}
\mu_{n\uparrow} &= \mu_{p\uparrow} + \mu_{e\uparrow} \\
&= \mu_{p\uparrow} + \mu_{e\downarrow} \\
&= \mu_{p\downarrow} + \mu_{e\uparrow}
\end{aligned} \tag{2}$$

$$\begin{aligned}
\mu_{n\downarrow} &= \mu_{p\downarrow} + \mu_{e\downarrow} \\
&= \mu_{p\downarrow} + \mu_{e\uparrow} \\
&= \mu_{p\uparrow} + \mu_{e\downarrow}
\end{aligned} \tag{3}$$

where up and down arrows indicate two directions of spin. In this case, the neutrinos which leave the system without delay, carry the spin signature of the processes. Of course, the neutrinos (anti-neutrinos) are assumed to be massless and left-handed (right-handed), i.e. they have definite helicity states. Now the general expression for the chemical potential for proton or neutron is given by

$$\begin{aligned}
\mu_p &= \epsilon_f^p + g_\omega \omega_0 + \frac{1}{2} g_\rho \rho_3^0 \\
\mu_n &= \epsilon_f^n + g_\omega \omega_0 - \frac{1}{2} g_\rho \rho_3^0
\end{aligned} \tag{4}$$

where the first terms in these equations are given by $\epsilon_f^i = (k_f^{i^2} + m^{*2})^{1/2}$, the neutron-proton effective single particle energy, $m^* = m - g_\sigma \sigma_0$ is the hadronic effective mass, g_i 's are the coupling constants, and

$$g_\sigma \sigma_0 = \left(\frac{g_\sigma}{m_\sigma} \right)^2 \sum_{i=n,p} \frac{g_i m^*}{2\pi^2} \sum_{\tau=\uparrow}^{\downarrow} \int_0^{k_{F_i}^{(\tau)}} \frac{k^2 dk}{(k^2 + m^{*2})^{1/2}} \tag{5}$$

Since there are sum over both iso-spin and spin, the explicit spin dependence of hadronic effective mass will not there, however there is an indirect spin dependence through the fermi momenta ($k_{F_i}^{(\tau)}$). The degeneracy parameter $g_i = 1$ for both $i = n$ and p . Now only the iso-spin 3-component of the iso-vector meson field has finite mean value, i.e., the neutral ρ meson (ρ_3^0). The Lorentz three-current part must vanish in the static ground state of matter, just as the vector meson ω^μ , leaving only ρ_3^0 . Since we are interested in spontaneous ferro-magnetic transition at low temperature (old neutron stars) in absence of external magnetic field, we have not considered the anomalous magnetic moments of the constituents. It is necessary to take into account the effect of anomalous magnetic moments to study the properties of nuclear matter in presence of strong magnetic field. From the above two sets of equations it is very easy to establish the following relations (since the spin dependence in chemical potentials come from the anomalous magnetic moments and is given by $s\delta_i B$, where $i = p, n$ or e , $s = +1$ for spin up and $s = -1$ for spin down cases, δ_i is related to the anomalous magnetic moment and finally B is the strength of magnetic field).

$$\mu_{n\uparrow} - \mu_{n\downarrow} = \mu_{p\uparrow} - \mu_{p\downarrow} = \mu_{e\uparrow} - \mu_{e\downarrow} = 0 \tag{6}$$

which further means

$$\epsilon_f^{p\uparrow} = \epsilon_f^{p\downarrow}; \quad \epsilon_f^{n\uparrow} = \epsilon_f^{n\downarrow}; \quad \epsilon_f^{e\uparrow} = \epsilon_f^{e\downarrow} \tag{7}$$

Hence it is extremely trivial to show from the definition of number density, that each species with up and down spins are equal in number, i.e. $n_{p\uparrow} = n_{p\downarrow}$, $n_{n\uparrow} = n_{n\downarrow}$, and $n_{e\uparrow} = n_{e\downarrow}$. *This is absolutely independent of the conventional models generally used to obtain the equation of states of NS matter.* Therefore, it is unlikely to have a spin polarized NS matter or a spontaneous ferromagnetic transition in NS, in which neutrinos flow out of the system as soon as they are created. Next we consider the case of a PNS, which is the very early stage of evolution of a typical NS, neutrinos remain trapped within the matter and it requires a finite duration of time (a few ms) to make the system neutrino free. In the case of PNS, since neutrinos are degenerate, their chemical potentials cannot be neglected in the dynamic equilibrium conditions as mentioned above (eqns.(2) and (3)). Therefore, the modified form of these two sets of equations are given by

$$\begin{aligned}
\mu_{n\uparrow} &= \mu_{p\uparrow} + \mu_{e\uparrow} - \mu_{\nu\downarrow} \\
&= \mu_{p\uparrow} + \mu_{e\downarrow} - \mu_{\nu\uparrow} \\
&= \mu_{p\downarrow} + \mu_{e\uparrow} - \mu_{\nu\uparrow}
\end{aligned} \tag{8}$$

$$\begin{aligned}
\mu_{n\downarrow} &= \mu_{p\downarrow} + \mu_{e\downarrow} - \mu_{\nu\uparrow} \\
&= \mu_{p\downarrow} + \mu_{e\uparrow} - \mu_{\nu\downarrow} \\
&= \mu_{p\uparrow} + \mu_{e\downarrow} - \mu_{\nu\downarrow}
\end{aligned} \tag{9}$$

From these two sets of conditions, it is very easy to show

$$\mu_{n\uparrow} - \mu_{n\downarrow} = \mu_{p\uparrow} - \mu_{p\downarrow} = \mu_{e\uparrow} - \mu_{e\downarrow} = \mu_{\nu\uparrow} - \mu_{\nu\downarrow} = S \quad (\text{say}) \tag{10}$$

where $|S| \neq 0$. This parameter determines the degree of polarization of the system, i.e. it controls the ferromagnetic transition of the system. In the case of unpolarized matter, $|S| \approx 0$, whereas for the spontaneously created ferromagnetic system, it is extremely large. From eqn.(10), it is also obvious that the magnitude of the factor S also significantly depends on the degree of polarization of both electrons and neutrinos / anti-neutrinos. In the case of electrons, the kinetic energy is found to be very high; therefore, the temperature of the system should be low enough to make them align. Since the main mechanism of NS/PNS cooling is by the emission of neutrinos, which in turn makes the neutrinos non-degenerate, and as a consequence the condition as shown above in eqn.(10) will no longer be valid. Instead, one has to consider eqn.(6), for which there will be no spontaneous magnetization, i.e., the process of alignment of electrons by reducing the temperature of the system through neutrino emission destroys the possibility of ferromagnetic transition. Therefore, it is very difficult to have a ferromagnetic phase transition even in PNS if the kinetic energy of the electronic part is high enough, which is sufficient to prevent spin alignment. If we concentrate on neutrino components, again there will be more serious problem than the electronic sector, when we consider the spin alignment of neutrinos. In the case, when the decaying neutron has spin up (down) and the recoiled proton has spin down (up), there will be some directional constraint on the motion of trapped neutrino / anti-neutrino (analogous to the famous ^{60}Co experiment) sum of electron and anti-neutrino spins should be $+1$ (-1). There will also be left-handed to right-handed and vice-versa flipping problem, unless the neutrinos / anti-neutrinos carry some non-zero mass.

Therefore, we conclude that in a very old neutron star, whose temperature is low enough and the neutrinos are non-degenerate, the spontaneous ferromagnetic transition is absolutely forbidden. On the other hand, in a PNS, where the neutrinos remain trapped within the matter for quite a long duration, are degenerate, both the electronic and neutrino components play major roles in ferromagnetic phase transition. The electron kinetic energy should be extremely low for their spin alignment, whereas, the neutrinos should have some finite non-zero mass (no definite helicity state) to have left-handed to right-handed spin flip and vice versa to make neutrinos / anti-neutrinos also spin aligned. Therefore, the final conclusion is that the neutrino mass has a significant role in spontaneous ferromagnetic phase transition in PNS. There will be no such transition if the neutrinos are mass less.

On the other hand for NS matter with arbitrary ISO-spin asymmetry parameter $x = (n_n - n_p)/n_B$, where $n_B = n_n + n_p$, the total baryon number density of the system, there could be a spontaneous ferromagnetic transition [10], however, the system will again become unpolarized as soon as β -equilibrium is reached, which we cannot avoid as the NS matter in β -equilibrium is the only stable configuration.

Therefore, if by some means a ferromagnetic transition occurs in PNS, then in the latter stage of evolution, when neutrinos become non-degenerate, there will be a spontaneous spin symmetry restoration. We may therefore conclude that the origin of residual magnetic field or the magnetic field of millisecond pulsars, which are believed to be very old neutron stars ($\sim 10^8\text{G}$), is not ferromagnetic transition. Now it is quite possible to have a crust of magnetic materials (e.g., iron) in such cold objects. To get an idea of what is the contribution of such magnetic materials on the bulk magnetic field of the object, we proceed in the following simplified manner. We give an estimate on the upper limit of polar magnetic field strength. If $\vec{m}$ is the total effective magnetic dipole moment of the crustal matter due to tiny atomic magnets (which are assumed to be aligned), we have the field strength at the pole

$$B_p = \frac{2 |\vec{m}|}{R^3} \tag{11}$$

where R is the radius of the star (assuming spherical). Assuming maximal projection of the total magnetic moment in unit of μ_B , the Bohr magneton, we have

$$B_p = \frac{8\pi}{3} g J \mu_B n \left[1 - \left(\frac{R'}{R} \right)^3 \right] \tag{12}$$

where R' is the thickness of the spherical crustal shell, g is the Lande g -factor and J is the total angular momentum of the electron and n is the density of tiny magnets. Then for iron $n = \rho/M$ and $gJ = 6$, where ρ is the density of crustal matter, which is assumed to be $10^{10} \text{ gm cm}^{-3}$ and M is the mass of tiny atomic magnets. Putting all these numerical numbers, we have seen that $B_p \approx 10 \text{ Tesla}$, which is of course too low. Again, this is the upper limit of polar magnetic field strength due to ferromagnetic transition of crustal magnetic material. Therefore, the ferromagnetic transition in the crustal matter can not produce magnetic field of the order $10^7 - 10^8 \text{ G}$.

- [1] S. Shapiro and S. Teukolsky, 1983, Black Holes, White Dwarfs, and Neutron Stars, John Wiley and Sons, Inc.
- [2] V. Ginzburg, 1964, Doklady Akad. Nauk, **156**, 43.
- [3] V. Ginzburg and L. Ozernoy, 1964, Sov. Phys. - ZhETF, **47**, 1030.
- [4] L. Woltjer, 1964, Astrophys. Jour., **140**, 1309
- [5] Ya. Zeldovich and I. Novikov, 1971, in Stars and Relativity, Univ. Chicago Press, Chicago.
- [6] M. Colpi, A. Posenti, S. Popov and F. Pizzolato, Spin and magnetism in old neutron stars, in "Physics of neutron stars interiors", Lecture notes in Physics, 578, Springer Verlag (2001), Eds. D. Blaschke, N.K. Glendenning and A. Sedrakian.
- [7] F. Pacini, Nature, (1967), **216**, 567.
- [8] T. Gold, Nature, (1968), **218**, 731.
- [9] J.P. Ostriker and J.E. Gunn, Astrophys. Jour., (1969), **157**, 1395.
- [10] Issac Vidaña and Ignazio Bombaci, nucl-th/0203061.

Acknowledgment: SC is thankful to the Department of Science and Technology, Govt. of India, for partial support of this work, Sanction number:SP/S2/K3/97(PRU). SC is also acknowledges Kamales Kar and Palas Pal of SINP for some useful discussions and comments.